



DEMOGRAPHIC RESEARCH

A peer-reviewed, open-access journal of population sciences

DEMOGRAPHIC RESEARCH

VOLUME 38, ARTICLE 32, PAGES 879–896

PUBLISHED 7 MARCH 2018

<http://www.demographic-research.org/Volumes/Vol38/32/>

DOI: 10.4054/DemRes.2018.38.32

Descriptive Finding

Estimating mortality from external causes using data from retrospective surveys: A validation study in Niakhar (Senegal)

Gilles Pison

Géraldine Duthé

Bruno Masquelier

Cheikh Sokhna

Almamy Malick Kante

Valérie Delaunay

Cheikh Tidiane Ndiaye

Stéphane Helleringer

Laetitia Douillot

© 2018 Gilles Pison et al.

This open-access work is published under the terms of the Creative Commons Attribution 3.0 Germany (CC BY 3.0 DE), which permits use, reproduction, and distribution in any medium, provided the original author(s) and source are given credit.

See <https://creativecommons.org/licenses/by/3.0/de/legalcode>.

Contents

| | | |
|-----|--|-----|
| 1 | Introduction | 880 |
| 2 | Methods | 882 |
| 2.1 | Study context and reference data | 882 |
| 2.2 | Validation study of survey data on deaths from external causes | 882 |
| 3 | Data analysis | 883 |
| 4 | Results | 885 |
| 5 | Discussion | 889 |
| 6 | Acknowledgments | 890 |
| | References | 891 |

Estimating mortality from external causes using data from retrospective surveys: A validation study in Niakhar (Senegal)

Gilles Pison¹

Géraldine Duthé⁶

Bruno Masquelier²

Cheikh Sokhna⁷

Almamy Malick Kante³

Valérie Delaunay⁵

Cheikh Tidiane Ndiaye⁴

Stéphane Helleringer³

Laetitia Douillot⁵

Abstract

BACKGROUND

In low- and middle-income countries (LMICs), data on causes of death is often inaccurate or incomplete. In this paper, we test whether adding a few questions about injuries and accidents to mortality questionnaires used in representative household surveys would yield accurate estimates of the extent of mortality due to external causes (accidents, homicides, or suicides).

METHODS

We conduct a validation study in Niakhar (Senegal), during which we compare reported survey data to high-quality prospective records of deaths collected by a health and demographic surveillance system (HDSS).

RESULTS

Survey respondents more frequently list the deaths of their adult siblings who die of external causes than the deaths of those who die from other causes. The specificity of

¹ Laboratoire d'Éco-anthropologie et ethnobiologie (UMR 7206 MNHN-CNRS-Paris Diderot), French Museum of Natural History, Paris, France and French Institute for Demographic Studies, Paris, France. Email: gilles.pison@mnhn.fr.

² French Institute for Demographic Studies, Paris, France and Centre for Demographic Research, Louvain University, Louvain-la-Neuve, Belgium.

³ Bloomberg School of Public Health, Johns Hopkins University, Baltimore, USA.

⁴ United Nations Population Fund, Moroni, Comores.

⁵ Unité mixte de recherches Lped, Institut de recherches pour le développement, Marseille, France.

⁶ French Institute for Demographic Studies, Paris, France.

⁷ Unité mixte de recherches Urmite, Institut de recherches pour le développement, Dakar, Sénégal.

survey data is high, but sensitivity is low. Among reported deaths, less than 60% of the deaths classified as due to external causes by the HDSS are also classified as such by survey respondents. Survey respondents better report deaths due to road-traffic accidents than deaths from suicides and homicides.

CONCLUSIONS

Asking questions about deaths resulting from injuries and accidents during surveys might help measure mortality from external causes in LMICs, but the resulting data displays systematic bias in a rural population of Senegal. Future studies should 1) investigate whether similar biases also apply in other settings and 2) test new methods to further improve the accuracy of survey data on mortality from external causes.

CONTRIBUTION

This study helps strengthen the monitoring of sustainable development targets in LMICs by validating a simple approach for the measurement of mortality from external causes.

1. Introduction

Reducing mortality due to external causes (e.g., accidents, suicides, or conflict-related deaths) figures prominently in the agenda for sustainable development. For example, halving global deaths and injuries from road-traffic accidents is one of the targets of the third Sustainable Development Goal (SDG) on ensuring healthy lives and promoting wellbeing. There is thus a strong need for accurate statistics on deaths due to external causes to 1) track progress toward the achievement of the SDGs and 2) evaluate the effectiveness of prevention initiatives and programs (Lagarde 2007).

Unfortunately, few LMICs have civil registration and vital statistics (CRVS) systems that permit monitoring mortality from external causes (AbouZahr et al. 2015; Mikkelsen et al. 2015; Phillips et al. 2014). In such countries, CRVS data is frequently incomplete and/or include large numbers of deaths attributed to unknown causes (Mikkelsen et al. 2015). Other data sources on injuries and accidents include police reports and hospital records, but both sources are affected by under-reporting (Abegaz et al. 2014; Sanghavi, Bhalla, and Das 2009; Sango et al. 2016). For example, large numbers of accident victims may never be transported to a hospital. CRVS systems often further break down during wars and other times of unrest. Data on conflict-related deaths thus comes from a number of incomplete sources, such as media reports (Hicks et al. 2011) or inventories of graveyards (Etcheson 2005). There is little agreement

between these sources (Carpenter, Fuller, and Roberts 2013), with estimates of a conflict's death toll routinely varying by orders of magnitude (Heuveline 2015).

Nevertheless, national levels and trends of mortality from external causes in LMICs are currently extrapolated from these partial data sources (Naghavi et al. 2017; Sanghavi, Bhalla, and Das 2009). The resulting estimates may be unreliable because they may omit some deaths, or inaccurately reclassify deaths attributed to unknown causes as injury-related (Bhalla and Harrison 2015). In studies of the global burden of disease (Naghavi et al. 2017), estimates of mortality from external causes are also frequently adjusted to ensure that they are consistent with other mortality estimates (e.g., deaths from infectious diseases). Errors in the measurement of other causes of death may further confound estimates of deaths due to external causes.

Improving the measurement of road-traffic mortality and other accidental deaths in LMICs thus requires collecting accurate and nationally representative data on deaths from external causes. There are ongoing efforts to strengthen CRVS systems (AbouZahr et al. 2015), but they mainly focus on birth registration and will have limited impact on our ability to measure causes of death in the next 15 years. Sample registration systems that include the collection of VA data are currently being established but only in a very limited number of countries (Kabadi et al. 2015; Mudenda et al. 2011).

We test whether mortality from external causes may be accurately measured during representative household surveys, such as the Demographic and Health Surveys (DHS). Surveys already constitute the primary data source on all-cause (Masquelier, Reniers, and Pison 2014) and maternal mortality (Zureick-Brown et al. 2013) for many LMICs. They measure mortality by asking respondents to report deaths among their relatives. Questions asking whether the deaths of these relatives are due to injuries, accidents, or conflicts have been asked in several surveys, such as DHS in Zimbabwe and Zambia, the WHO's World Health Survey (Obermeyer, Murray, and Gakidou 2008), and postconflict surveys (Hagopian et al. 2013; Lafta et al. 2015). A few studies have compared estimates of conflict-related deaths derived from such surveys to estimates from other sources (Obermeyer, Murray, and Gakidou 2008). But the accuracy of survey data in recording other external causes of death has not been evaluated.

2. Methods

2.1 Study context and reference data

We conduct a validation study of survey data on mortality from external causes. We use data from the Niakhar health and demographic surveillance system (HDSS) in Senegal (West Africa) as reference. HDSS are populations in which vital events (e.g., births, deaths) are recorded continuously during periodic household visits (Pison 2005). Each recorded death is also investigated by verbal autopsy (VA), i.e., a postmortem interview with caregivers of the deceased. VA data is highly accurate in identifying deaths from external causes (Baiden et al. 2007; Chandramohan et al. 1994, 1998).

In Niakhar, after a baseline census, data on births, deaths, marriages, and migrations are collected from household informants during regular household visits (Delaunay et al. 2013). Each recorded death is investigated using VA. Two clinicians then review VA data and attribute a probable cause using codes of the international classification of diseases. The most frequent external causes of death recorded in the Niakhar HDSS are falls, suicides, road accidents, fire, and burning (Guyavarch et al. 2010).

The Niakhar HDSS constitutes a high-quality reference dataset on mortality from external causes, but it has limitations. First, some external causes of death (e.g., war deaths) are not represented in this dataset. If such deaths are more (or less) precisely reported during surveys than other accidents and injuries, our study may misrepresent the accuracy of survey-based estimates in other settings. Second, despite the high accuracy of VA data about mortality from external causes, HDSS data does not constitute a gold standard. It may still misclassify some external causes of death as nonexternal, and vice versa.

2.2 Validation study of survey data on deaths from external causes

In 2013, we conducted a survey of adult mortality among the Niakhar HDSS population. Our goal was to compare survey reports of adult deaths among the siblings of a respondent to prospective HDSS records of the survival of these siblings.

We selected a stratified sample of (a) sibships in which one sibling (male or female) died between 15 and 59 years old ($n=592$) and (b) other sibships in which no adult death was recorded by the HDSS ($n=500$). We then selected at random up to two surviving members of each sibship for a survey interview. Respondents included both men and women aged 15–59 years old. Study interviewers were unaware of the composition of respondents' sibships.

In total, 1,189 respondents were asked to list their maternal siblings by birth order and report their survival statuses, ages, and, if deceased, ages at death and dates of death. For siblings who had died at ages 12 and older, respondents also answered a) whether their siblings died after being injured, b) whether the injuries were due to accidents, suicides, homicides, wars, or natural disasters, and c) the circumstances that lead to the injuries (e.g., whether they were due to collisions with motor vehicles).

We test two adult mortality questionnaires: (1) the standard DHS questionnaire and (2) the siblings' survival calendar (SSC). The SSC incorporates supplementary interviewing techniques to limit omissions of siblings and uses an event history calendar to improve reports of dates and ages. The SSC improves the completeness and accuracy of data on all-cause adult mortality and maternal mortality (Helleringer et al. 2014; Helleringer et al. 2015). Exploratory analyses suggest that the SSC might also improve some aspects of the reporting of deaths from external causes, but the differences we observe are only marginally significant (see Discussion). In this paper, we thus pool data from the two questionnaires.

3. Data analysis

Two types of reporting errors affect survey data on deaths from external causes. First, selective omissions happen when the likelihood of reporting siblings' deaths depends on their causes of death. For example, a respondent may list their deceased brother if he died during a car accident but not if he died from HIV/AIDS. Second, misclassifications occur when respondents correctly list the deaths of their siblings but misstate the circumstances of their deaths. For example, a respondent may mistakenly state that their sister, who died after falling from a window, did not die from an injury or accident. Selective omissions and misclassifications may bias estimates of the proportion of adult deaths that are attributable to external causes (PEC hereafter). For example, if respondents do not list 10% of their siblings whose deaths are due to external causes vs. 20% of their siblings whose deaths are due to other causes, the PEC will be overestimated.

We focused on sibships in which a man or a woman died at age 12 years or above in the 15 years prior to the survey, i.e., the time frame commonly used in Global Burden of Disease studies (Wang et al. 2012). We first describe the characteristics of deaths recorded by the HDSS, and we compare the characteristics of deaths from external vs. other causes. These characteristics include the size of the sibship of the deceased, gender, age at death, and the time elapsed since the death. For deaths from external causes, we describe the circumstances of the deaths according to the HDSS datasets.

Second, we measure the extent of selective omissions, i.e., whether the probability of listing a deceased sibling during the survey varies by cause of death (external vs. other). Third, we measure misclassifications in survey reports of deaths from external causes. We define the sensitivity of survey data as the proportions of respondents correctly stating the cause of reported deaths among respondents in sibships with deaths from external causes. We define the specificity of survey data as the proportions of respondents correctly stating the cause of reported deaths among respondents in sibships with only deaths from nonexternal causes. We calculate 95% confidence intervals around each of these estimates after a logit transform (so that the endpoints lie between 0 and 1). We use Student's t-distribution (with degrees of freedom equal to the number of observations minus 1) to determine the critical values of the confidence intervals. We obtain estimates of each proportion (as well as its standard error and confidence interval) using the `svy` command in Stata 14. In this setup, we consider the selection of sibships as our first sampling stage and the selection of respondents within sibships as our final sampling stage. This allows adjusting standard errors and confidence intervals for the clustering of survey reports in sibships with several respondents.

Fourth, we use estimates of these parameters to evaluate bias in survey estimates of the PEC using the following formula:

$$\widehat{PEC} = \pi \times R_{EC} \times Se + (1 - \pi) \times R_O \times Sp, \quad (1)$$

in which \widehat{PEC} is the survey estimate of the PEC, π is the true value of this proportion, R_{EC} is the probability that respondents will list their sibling who died from external causes, R_O is a similar probability for siblings who died from other causes, and Se and Sp are the sensitivity and specificity of survey data in classifying death from external causes, respectively. We explore how much variations in each of the reporting parameters (R_{EC} , R_O , Se , and Sp) in Equation (1) affect bias in survey estimates of the PEC. We thus let each of these parameters vary between the lower and upper bounds of its confidence interval, and we recalculate the \widehat{PEC} for each level of that range.

We investigate the joint effects of the four reporting parameters on estimates of the PEC. We let π vary between 0 and 0.3, in which the upper bound corresponds to one of the highest PEC observed among adult residents of HDSS, i.e., Mlomp in Southern Senegal (Guyavarch *et al.* 2010). For each value of π varying between 0 and 0.3, we generate 500 estimates of the \widehat{PEC} through draws from the distributions of the reporting parameters in conjunction with Equation (1). We calculate the median \widehat{PEC} for each level of π . We also calculate the 2.5th and 97.5th percentiles of the distribution of PEC estimates to construct a 95% confidence interval around this median value.

Finally, to explore whether some external deaths may be more likely to be omitted or misclassified during surveys, we compare the circumstances of deaths recorded by the HDSS between ‘false negatives’ and ‘true positives.’ False negatives are instances in which survey respondents do not classify deaths as due to external causes when the HDSS does so, whereas true positives are instances in which both respondents and the HDSS attribute the deaths to external causes.

4. Results

There are 468 deaths of persons aged 12 years or older during the 15 years before the survey. Twenty-five of these deaths (5.5 %) could be attributed to external causes on the basis of HDSS data (Table 1). The PEC is higher for males than for females (6.7% vs. 2.6%).

Table 1: Characteristics of deaths recorded by the Niakhar HDSS during the 15 years prior to the survey and included in the validation study

| | External causes (N=25) | Other causes (N=443) |
|-------------------------------|---------------------------|-------------------------|
| Sibship size | | |
| 1–4 maternal siblings | 6 (4.4) | 130 (95.6) |
| 5+ maternal siblings | 19 (5.7) | 313 (94.3) |
| Gender | | |
| Male | 21 (6.7) | 291 (93.3) |
| Female | 4 (2.6) | 152 (97.4) |
| Age at death | | |
| <25 years old | 10 (5.9) | 159 (94.1) |
| 25–34 years old | 8 (6.5) | 115 (93.5) |
| 35–44 years old | 5 (5.8) | 82 (94.3) |
| ≥45 years old | 2 (2.3) | 87 (97.7) |
| Time since death | | |
| ≤5 years | 11 (6.1) | 169 (93.9) |
| 6–10 years | 7 (4.2) | 160 (95.8) |
| 11–15 years | 7 (5.8) | 114 (94.2) |
| Circumstances of death | | |
| Falls/drowning | 4 (16.0) | – |
| Road-traffic accidents | 5 (20.0) | – |
| Homicide/suicide | 4 (16.0) | – |
| Other accidents/unknown | 12 (48.0) | – |

Note: Figures in parentheses are row percentages.

The survey data is affected by selective omissions of deaths (Table 2). Among respondents whose siblings died of external causes according to HDSS, 34 out of 35 report a death at ages 12 or older (97.1%). The proportion is lower (497 out of 567, 87.7%) among respondents whose siblings died of nonexternal causes ($p=0.087$).

Table 2: Selective omissions of adult deaths during survey interviews by sibship composition

| | Sibships with adult deaths from external causes | | Sibships with adult deaths from other causes | | p-value |
|-------------|---|----------------------|--|----------------------|---------|
| | Reported/expected | % (95% CI) | Reported/expected | % (95% CI) | |
| All deaths | 34/35 | 97.1 (82.8, 99.6) | 497/567 | 87.7 (84.3, 90.4) | 0.087 |
| Male deaths | 31/32 | 96.9 (81.4, 99.6) | 334/383 | 87.2 (83.0, 90.5) | 0.102 |

Note: 'Reported' refers to the number of respondents who report an adult death among their siblings during the past 15 years; 'expected' refers to the number of respondents who are members of sibships in which there are at least one death due to injuries or one death due to other causes in the past 15 years; p-values test the null hypothesis of no difference in proportion of respondents reporting an adult death during the past 15 years between sibships with an injury death and sibships with only noninjury deaths. This p-value is based on a χ^2 test. Standard errors are adjusted for the clustering of respondents within sibships using the svy command in Stata.

The sensitivity of survey data is low (Table 3): In sibships with a death attributed to external causes by HDSS, 20 out of 34 (58.8%) survey respondents correctly classify the reported death as due to external causes. The specificity of survey data is high: In sibships with only deaths from other causes according to HDSS, 471 out of 497 respondents correctly classify the reported death (94.8%).


Table 3: Sensitivity/specificity of reports of adult deaths collected during survey interviews

| | Sensitivity | | Specificity | | p-value |
|-------------|-------------------|----------------------|-------------------|----------------------|---------|
| | Correct/ reported | % (95% CI) | Correct/ reported | % (95% CI) | |
| All deaths | 20/34 | 58.8 (41.0, 74.6) | 471/497 | 94.8 (92.0, 96.6) | <0.001 |
| Male deaths | 19/31 | 61.3 (43.2, 76.7) | 318/334 | 95.2 (91.5, 97.4) | <0.001 |

Note: 'Sensitivity' refers to the proportion of deaths classified as due to injuries by the HDSS dataset that are also classified as due to injuries during the survey. Specificity refers to the proportion of deaths classified as due to other causes of death (i.e., noninjuries) by the HDSS dataset that are also classified as such during the survey. P-values test the null hypothesis of no difference between sensitivity and specificity. This p-value is based on a χ^2 test. Standard errors are adjusted for the clustering of respondents within sibships using the svy command in Stata.

Selective omissions during surveys as measured in this study (Figures 1a and 1b) have limited impact on bias in PEC estimates. But the sensitivity/specificity of survey data (Figures 1c and 1d) could modify our inferences about the PEC. For example, in settings with low prevalence of deaths from external causes, survey estimates of the PEC are significantly affected by small variations in the specificity of survey data, but this is not the case in contexts with high prevalence of such deaths (Figure 1d).

Figure 1: Effects of reporting parameters on bias in survey estimates of the PEC




Notes: In these figures, the bias ratio is derived from calculations using Equation (1). Specifically, it is defined as (1) the difference between the PEC calculated for a specific value of the parameter of interest (within its confidence interval) and the true value of π (either 0.05 or 0.25) divided by (2) the difference between the PEC calculated for the point estimate of that parameter and the true value of π . A bias ratio > 1 indicates that the bias is larger for that value of the parameter than it is for the point estimate of that parameter. A value less than 1 indicates the opposite. The dotted vertical line represents the estimate of this parameter, as reported in Tables 2 and 3. The horizontal dotted line represents a bias ratio of 1.

Based on estimates of R_{EC} , R_O , Se , Sp , and their sampling variability, the direction of bias in survey estimates of the PEC (Equation 1) likely varies across populations.

When the true PEC is below approximately 13%, survey data overestimate the PEC (Figure 2), but in populations in which the true PEC is above 13%, survey data underestimate the PEC.

Figure 2: Joint effects of reporting errors on survey estimates of the PEC



Note: The dotted black line represents the 45-degree line, in which the reported proportion of deaths due to external causes is equal to the hypothesized true proportion of deaths due to external causes in the population. The solid line represents the expected survey estimate based on Equation (1) and estimates of the reporting parameters. Its confidence interval is calculated from 500 draws from the distributions of these reporting parameters. The distance between the dotted line and the expected survey estimate (solid line) measures the extent of bias in survey estimates of the PEC. For reference, the solid red line at 0.055 represents the proportion of deaths due to external causes in the study sample according to the HDSS. Other vertical lines represent estimates of the PEC among adults (aged 15–49 years old) in other HDSS in other LMICs. The estimate from Kersa (Ethiopia) is described in the following references (Ashenafi et al. 2017; Assefa et al. 2016). The estimate from Dande (Angola) is described in the following references (Rosario et al. 2016; Rosario et al. 2017). The estimate from Mlomp (Senegal) is described in the following reference (Guyavarch et al. 2010).

Four out of six survey reports in sibships in which adult deaths are classified as homicides or suicides by HDSS are false negatives, i.e., they are reported as due to nonexternal causes by survey respondents (Table 4). All the survey reports in sibships in which adult deaths are classified as road-traffic accidents by HDSS and four out of

five reports in sibships in which adult deaths are due to falls or drownings are true positives, i.e., they are correctly classified as due to external causes by survey respondents.

Table 4: Causes of death among deaths from external causes by reporting statuses during survey

| Circumstances of death | All | By survey reports | | p-value |
|-------------------------|-----------|-------------------|----------------|---------|
| | | False negatives | True positives | |
| Falls/drowning | 5 (14.3) | 1 (6.7) | 4 (20.0) | 0.069 |
| Road-traffic accidents | 5 (14.3) | 0 (0.0) | 5 (25.0) | |
| Homicide/suicide | 6 (17.1) | 4 (26.7) | 2 (10.0) | |
| Other accidents/unknown | 19 (54.3) | 10 (66.7) | 9 (45.0) | |

Note: False negatives are deaths that are classified as injury-related by the HDSS but are not classified as such during the survey. True positives are deaths that are classified as injury-related both by the HDSS and the survey. The difference in distribution of causes of death between false negatives and true positives is significant at $p < 0.1$ level according to Fisher's exact test.

5. Discussion

Adding questions about injuries and accidents to representative surveys conducted in LMICs might help measure mortality from external causes, but the resulting data displays systematic biases. In Senegal, survey respondents are more likely to list the deaths of their adult siblings if they die from external causes. Their reports of the circumstances of deaths have low sensitivity but high specificity in identifying deaths from external causes.

As a result, the accuracy of survey data in estimating mortality from external causes likely varies across populations. In populations in which such causes account for less than 13% of deaths, survey data yields estimates of the proportion of deaths due to external causes that are too high. In populations in which deaths from external causes are more common (>13%), survey data under-estimates this proportion. The accuracy of survey-based estimates may further depend on the types of injuries and accidents that are most prevalent in a population. Survey respondents accurately report road-traffic accidents, falls, and drowning but under-report or misclassify stigmatized causes of death, such as suicides and homicides.

Our study has limitations. First, due to small sample sizes, our measures of data accuracy are imprecise. We also cannot explore the correlates of reporting errors in survey data on deaths from external causes. Second, the wording of some of the survey questions may be too narrow. In particular, we ask respondents whether their siblings died following injuries, even though some deaths from external causes may not be

accompanied by injuries. Third, the Niakhar HDSS population may not be representative of other populations in LMICs. Some external causes of death are not represented in this setting. In part because it is a rural population with limited road-traffic, its inhabitants may also better recall car accidents than inhabitants of urban areas where such events are more common. Finally, because of long-term HDSS activities, Niakhar residents may be better informed about causes of death than members of other populations.

Methodological research is needed in several areas to improve the accuracy of survey data on mortality from external causes. New questions should be developed that better capture the diversity of accidents and injuries. More confidential interview techniques should be tested to improve the sensitivity of survey data, in particular as it relates to the reporting of stigmatized causes of death, such as suicides and homicides. Techniques such as audio computer-assisted self-interviewing (Mensch et al. 2008) or voting methods, for example, have helped reduce social desirability biases in the reporting of other sensitive behaviors (e.g., sexual risk taking). They may have similar effects on the reporting of under-reported violent deaths.

Strategies to further improve the specificity of survey data are also needed, since estimates of the PEC are affected by this parameter in settings in which mortality from external causes is low. The siblings' survival calendar tested as part of this study (Helleringer et al. 2014) might represent one such strategy. In exploratory analyses, it displayed higher specificity than the DHS questionnaire (97.2 vs. 92.2%, $p = 0.06$). Such improved survey instruments should be validated in additional LMIC populations. This will permit establishing whether results from this study are generalizable beyond the Niakhar HDSS and/or better describe heterogeneity in reporting behaviors across LMICs. Finally, new statistical models (e.g., Bayesian models) that incorporate prior information on data accuracy should be adopted for the estimation of mortality from external causes.

6. Acknowledgments

This work was supported by the Eunice Kennedy Shriver National Institute of Child Health and Human Development [R03-HD071117] and the French Agence Nationale de la Recherche [ANR-11-BSH1-0007]. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The authors declare that no competing interests exist.

References

- Abegaz, T., Berhane, Y., Worku, A., Assrat, A., and Assefa, A. (2014). Road-traffic deaths and injuries are under-reported in Ethiopia: A capture–recapture method. *PLoS One* 9(7): e103001. doi:10.1371/journal.pone.0103001.
- AbouZahr, C., de Savigny, D., Mikkelsen, L., Setel, P.W., Lozano, R., Nichols, E., Notzon, F., and Lopez, A.D. (2015). Civil registration and vital statistics: Progress in the data revolution for counting and accountability. *Lancet* 386(10001): 1373–1385. doi:10.1016/S0140-6736(15)60173-8.
- Ashenafi, W., Eshetu, F., Assefa, N., Oljira, L., Dedefo, M., Zelalem, D., Baraki, N., and Demena, M. (2017). Trend and causes of adult mortality in Kersa Health and Demographic Surveillance System (Kersa HDSS), Eastern Ethiopia: Verbal autopsy method. *Population Health Metrics* 15(1): 22. doi:10.1186/s12963-017-0144-2.
- Assefa, N., Oljira, L., Baraki, N., Demena, M., Zelalem, D., Ashenafi, W., and Dedefo, M. (2016). HDSS profile: The Kersa Health and Demographic Surveillance System. *International Journal of Epidemiology* 45(1): 94–101. doi:10.1093/ije/dyv284.
- Baiden, F., Bawah, A., Biai, S., Binka, F., Boerma, T., Byass, P., Chandramohan, D., Chatterji, S., Engmann, C., Greet, D., Jakob, R., Kahn, K., Kunii, O., Lopez, A.D., Murray, C.J., Nahlen, B., Rao, C., Sankoh, O., Setel, P.W., Shibuya, K., Soleman, N., Wright, L., and Yang, G. (2007). Setting international standards for verbal autopsy. *Bulletin of the World Health Organization* 85(8): 570–571. doi:10.2471/BLT.07.043745.
- Bhalla, K. and Harrison, J.E. (2015). GBD–2010 overestimates deaths from road injuries in OECD countries: New methods perform poorly. *International Journal of Epidemiology* 44(5): 1648–1656. doi:10.1093/ije/dyv019.
- Carpenter, D., Fuller, T., and Roberts, L. (2013). WikiLeaks and Iraq Body Count: The sum of parts may not add up to the whole: A comparison of two tallies of Iraqi civilian deaths. *Prehospital and Disaster Medicine* 28(3): 223–229. doi:10.1017/S1049023X13000113.
- Chandramohan, D., Maude, G.H., Rodrigues, L.C., and Hayes, R.J. (1994). Verbal autopsies for adult deaths: Issues in their development and validation. *International Journal of Epidemiology* 23(2): 213–222. doi:10.1093/ije/23.2.213.

- Chandramohan, D., Maude, G.H., Rodrigues, L.C., and Hayes, R.J. (1998). Verbal autopsies for adult deaths: Their development and validation in a multicentre study. *Tropical Medicine and International Health* 3(6): 436–446. doi:10.1046/j.1365-3156.1998.00255.x.
- Delaunay, V., Douillot, L., Diallo, A., Dione, D., Trape, J.F., Medianikov, O., Raoult, D., and Sokhna, C. (2013). Profile: The Niakhar Health and Demographic Surveillance System. *International Journal of Epidemiology* 42(4): 1002–1011. doi:10.1093/ije/dyt100.
- Etcheson, C. (2005). *After the killing fields: Lessons from the Cambodian genocide*. Westport: Praeger.
- Guyavarch, E., Pison, G., Duthé, G., Marra, A., and Chippaux, J.-P. (2010). Mortality due to external causes in three rural areas of Senegal. *European Journal of Population/Revue européenne de Démographie* 26(4): 483–505. doi:10.1007/s10680-010-9213-y.
- Hagopian, A., Flaxman, A.D., Takaro, T.K., Esa Al Shatari, S.A., Rajaratnam, J., Becker, S., Levin-Rector, A., Galway, L., Hadi Al-Yasseri, B.J., Weiss, W.M., Murray, C.J., and Burnham, G. (2013). Mortality in Iraq associated with the 2003–2011 war and occupation: Findings from a national cluster sample survey by the University Collaborative Iraq Mortality Study. *PLoS Medicine* 10(10): e1001533. doi:10.1371/journal.pmed.1001533.
- Helleringer, S., Pison, G., Masquelier, B., Kante, A.M., Douillot, L., Duthé, G., Sokhna, C., and Delaunay, V. (2014). Improving the quality of adult mortality data collected in demographic surveys: Validation study of a new siblings' survival questionnaire in Niakhar, Senegal. *PLoS Medicine* 11(5). doi:10.1371/journal.pmed.1001652.
- Helleringer, S., Pison, G., Masquelier, B., Kante, A.M., Douillot, L., Ndiaye, C.T., Duthé, G., Sokhna, C., and Delaunay, V. (2015). Improving survey data on pregnancy-related deaths in low- and middle-income countries: A validation study in Senegal. *Tropical Medicine and International Health* 20(11): 1415–1423. doi:10.1111/tmi.12583.
- Heuveline, P. (2015). The boundaries of genocide: Quantifying the uncertainty of the death toll during the Pol Pot regime in Cambodia, 1975–1979. *Population Studies* 69(2): 201–218. doi:10.1080/00324728.2015.1045546.

- Hicks, M.H., Dardagan, H., Guerrero Serdan, G., Bagnall, P.M., Sloboda, J.A., and Spagat, M. (2011). Violent deaths of Iraqi civilians, 2003–2008: Analysis by perpetrator, weapon, time, and location. *PLoS Medicine* 8(2): e1000415. doi:10.1371/journal.pmed.1000415.
- Kabadi, G.S., Geubbels, E., Lyatuu, I., Smithson, P., Amaro, R., Meku, S., Schellenberg, J.A., and Masanja, H. (2015). Data resource profile: The sentinel panel of districts: Tanzania's national platform for health impact evaluation. *International Journal of Epidemiology* 44(1): 79–86. doi:10.1093/ije/dyu223.
- Lafta, R., Al-Shatari, S., Cherewick, M., Galway, L., Mock, C., Hagopian, A., Flaxman, A., Takaro, T., Greer, A., Kushner, A., and Burnham, G. (2015). Injuries, death, and disability associated with 11 years of conflict in Baghdad, Iraq: A randomized household cluster survey. *PLoS One* 10(8): e0131834. doi:10.1371/journal.pone.0131834.
- Lagarde, E. (2007). Road traffic injury is an escalating burden in Africa and deserves proportionate research efforts. *PLoS Medicine* 4(6): e170. doi:10.1371/journal.pmed.0040170.
- Masquelier, B., Reniers, G., and Pison, G. (2014). Divergences in trends in child and adult mortality in sub-Saharan Africa: Survey evidence on the survival of children and siblings. *Population Studies* 68(2): 161–177. doi:10.1080/00324728.2013.856458.
- Mensch, B.S., Hewett, P.C., Gregory, R., and Helleringer, S. (2008). Sexual behavior and STI/HIV status among adolescents in rural Malawi: An evaluation of the effect of interview mode on reporting. *Studies in Family Planning* 39(4): 321–334. doi:10.1111/j.1728-4465.2008.00178.x.
- Mikkelsen, L., Phillips, D.E., AbouZahr, C., Setel, P.W., de Savigny, D., Lozano, R., and Lopez, A.D. (2015). A global assessment of civil registration and vital statistics systems: Monitoring data quality and progress. *Lancet* 386(10001): 1395–1406. doi:10.1016/S0140-6736(15)60171-4.
- Mudenda, S.S., Kamocha, S., Mswia, R., Conkling, M., Sikanyiti, P., Potter, D., Mayaka, W.C., and Marx, M.A. (2011). Feasibility of using a World Health Organization-standard methodology for sample vital registration with verbal autopsy (SAVVY) to report leading causes of death in Zambia: Results of a pilot in four provinces, 2010. *Population Health Metrics* 9: 40. doi:10.1186/1478-7954-9-40.

- Naghavi, M., Abajobir, A.A., Abbafati, C., Abbas, K.M., Abd-Allah, F., Abera, S.F., et al. (2017). Global, regional, and national age–sex specific mortality for 264 causes of death, 1980–2016: A systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 390(10100): 1151–1210. doi:10.1016/S0140-6736(17)32152-9.
- Obermeyer, Z., Murray, C.J., and Gakidou, E. (2008). Fifty years of violent war deaths from Vietnam to Bosnia: Analysis of data from the World Health Survey Programme. *British Medical Journal* 336(7659): 1482–1486. doi:10.1136/bmj.a137.
- Phillips, D.E., Lozano, R., Naghavi, M., Atkinson, C., Gonzalez-Medina, D., Mikkelsen, L., Murray, C.J., and Lopez, A.D. (2014). A composite metric for assessing data on mortality and causes of death: The vital statistics performance index. *Population Health Metrics* 12: 14. doi:10.1186/1478-7954-12-14.
- Pison, G. (2005). Population observatories as sources of information on mortality in developing countries. *Demographic Research* 13(13): 301–334. doi:10.4054/DemRes.2005.13.13.
- Rosario, E.V., Costa, D., Timoteo, L., Rodrigues, A.A., Varanda, J., Nery, S.V., and Brito, M. (2016). Main causes of death in Dande, Angola: Results from verbal autopsies of deaths occurring during 2009–2012. *BMC Public Health* 16: 719. doi:10.1186/s12889-016-3365-6.
- Rosario, E.V.N., Costa, D., Francisco, D., and Brito, M. (2017). HDSS profile: The Dande Health And Demographic Surveillance System (Dande HDSS, Angola). *International Journal of Epidemiology* 46(4): 1094–1094g. doi:10.1093/ije/dyx072.
- Sanghavi, P., Bhalla, K., and Das, V. (2009). Fire-related deaths in India in 2001: A retrospective analysis of data. *Lancet* 373(9671): 1282–1288. doi:10.1016/S0140-6736(09)60235-X.
- Sango, H.A., Testa, J., Meda, N., Contrand, B., Traore, M.S., Staccini, P., and Lagarde, E. (2016). Mortality and morbidity of urban road traffic crashes in Africa: Capture–recapture estimates in Bamako, Mali, 2012. *PLoS One* 11(2): e0149070. doi:10.1371/journal.pone.0149070.

- Wang, H., Dwyer-Lindgren, L., Lofgren, K.T., Rajaratnam, J.K., Marcus, J.R., Levin-Rector, A., Levitz, C.E., Lopez, A.D., and Murray, C.J. (2012). Age-specific and sex-specific mortality in 187 countries, 1970–2010: A systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 380(9859): 2071–2094. doi:10.1016/S0140-6736(12)61719-X.
- Zureick-Brown, S., Newby, H., Chou, D., Mizoguchi, N., Say, L., Suzuki, E., and Wilmoth, J. (2013). Understanding global trends in maternal mortality. *International Perspectives on Sexual and Reproductive Health* 39(1): 32–41. doi:10.1363/3903213.

