

DEMOGRAPHIC RESEARCH

A peer-reviewed, open-access journal of population sciences

DEMOGRAPHIC RESEARCH

**VOLUME 44, ARTICLE 19, PAGES 443–458
PUBLISHED 9 MARCH 2021**

<https://www.demographic-research.org/Volumes/Vol44/19/>

DOI: 10.4054/DemRes.2021.44.19

Research Material

The centered ternary balance scheme: A technique to visualize surfaces of unbalanced three-part compositions

Jonas Schöley

This article is part of the Special Collection on “Data Visualization,” organized by Guest Editors Tim Riffe, Sebastian Klüsener, and Nikola Sander.

© 2021 Jonas Schöley.

This open-access work is published under the terms of the Creative Commons Attribution 3.0 Germany (CC BY 3.0 DE), which permits use, reproduction, and distribution in any medium, provided the original author(s) and source are given credit.

See <https://creativecommons.org/licenses/by/3.0/de/legalcode>.

Contents

1	Ternary diagrams and color schemes	444
2	The challenge of unbalanced compositions	445
3	Perturbation and centering of compositional data	447
4	The centered ternary balance scheme	448
5	Discussion	450
6	Acknowledgments	454
	References	455

The centered ternary balance scheme: A technique to visualize surfaces of unbalanced three-part compositions

Jonas Schöley¹

Abstract

BACKGROUND

The ternary balance scheme is a visualization technique that encodes three-part compositions as a mixture of three primary colors. The technique works best if the compositional data are well spread out across the domain but fails to show structure in very unbalanced data.

OBJECTIVE

I extend the ternary balance scheme such that it can be utilized to show variation in unbalanced compositional surfaces.

METHODS

By reprojecting an unbalanced compositional data set around its center of location and visualizing the transformed data with a standard ternary balance scheme, the internal variation of the data becomes visible. An appropriate centering operation has been defined within the field of compositional data analysis.

RESULTS

Using Europe's regional workforce structure by economic sector as an example, I have demonstrated the utility of the centered ternary balance scheme in visualizing variation across unbalanced compositional surfaces.

CONTRIBUTION

I have proposed a technique to visualize the internal variation in surfaces of highly unbalanced compositional data and implemented it in the *tricolore* R package.

¹ Interdisciplinary Centre on Population Dynamics, University of Southern Denmark.
ORCID: 0000-0002-3340-8518.
Email: jschoeley@sdu.dk.


1. Ternary diagrams and color schemes

When it comes to proportions, the number 3 is quite significant: the share of people working in the primary vs. secondary vs. tertiary sector, the proportion of total population change explained by migration vs. fertility vs. mortality, the relative population numbers in young age vs. working age vs. retirement age, the share of a cohort attaining primary vs. secondary vs. tertiary education degrees, the relative number of deaths due to prematurity vs. accidents vs. old age, the share of papers accepted as is vs. revised vs. rejected and so on: Three-part proportions of a whole, i.e., ternary compositions, are a type of data that is both ubiquitous and idiosyncratic enough as to warrant particular attention when it comes to presentation. The ternary diagram and its use throughout the sciences stand as a manifestation of this view.

Variably referred to as a de Finetti, simplex, or triangle plot, the ternary diagram is based upon a coordinate system that maps each point within an equilateral triangle to a unique three-part composition and thus has found use wherever the problem domain spans three parts of a whole. The diagram emerged during the 18th century as a means of illustrating relative mixtures of primary colors (Howarth 1996). It was subsequently adopted as the standard method to depict phase transitions in three-component alloys (Bancroft 1897), the genotype composition of a population (De Finetti 1926), soil composition (Davis and Hammond 1927), and the potential for flammability given different mixtures of three gases (Zabetakis 1965). In the social sciences, ternary diagrams depict population compositions along demographic characteristics, with an early example appearing in the USSR's first census report showing the distribution of workers across labor market segments in various regions (Kvitkin 1932).

Wherever three-part compositions are available by geographical region or other pairs of ordered attributes, such as cohort and age, one faces the challenge of visualizing ternary compositions on a surface such as the surface of the Earth or the period-age Lexis surface. The ternary balance scheme (Brewer 1994) is a color scale suited to that task. The technique encodes the relative shares among three parts as a mixture of three primary colors. Figure 1b shows the proportions of people with either “lower secondary or less,” “secondary,” or “tertiary” educational attainment by European region in 2016. Lower degrees are mapped to yellow, secondary to cyan, and tertiary to magenta. The more pronounced the yellow in a region, the higher the share of people with lower education. The same logic applies to the two other education categories. The more greyish a region is colored, the more balanced the three proportions are, with a perfect grey signifying an equal share of people in all three education categories. A ternary diagram is used as a color key (see Figure 1a) and doubles as a visualization of the distribution of data marginalized over the geographical surface.

Figure 1: Demonstration of the ternary balance scheme showing the composition of educational attainment by region in Europe, 2016. Data by Eurostat.




Published examples of the ternary balance scheme include maps of population compositions by political alignment, education, and workforce status (Dorling 2012; Graetz et al. 2019; Brewer 1994), geological maps of soil composition (Metternicht and Stott 2003), maps of Arctic sea ice coverage by type (Denil 2015), and land cover compositions by type of forest (Pirzamanbein, Poska, and Lindström 2020; Steidinger et al. 2019). Schöley and Willekens (2017) employed the scheme to visualize the distribution of deaths by cause among the French population on a period by age (Lexis) surface.

2. The challenge of unbalanced compositions

While the ternary balance scheme allows for incredibly dense yet clear visualizations of well-spread-out three-part compositions, the technique is less informative when used with highly unbalanced data. Figure 2a shows the regional workforce composition in Europe as of 2016. The map is almost monochromatic, with the intense magenta signifying a working population concentrated in the tertiary (services) sector. Regions in Turkey and Eastern Europe show a somewhat higher concentration of workers in the primary (production) sector, but overall there is little variation with regard to the visual reference point, i.e., the grey-point marking perfectly balanced proportions.

Figure 2: Demonstration of the centered ternary balance scheme in comparison with the non-centered scheme showing the workforce composition by region in Europe, 2016. Data by Eurostat.



A remedy for analyzing data that show little variation in relation to some reference point is to adjust the point of reference. Figure 2b yet again shows the European regional workforce composition, but the data have been reprojected so that the center of the compositional point cloud, the average European workforce composition, coincides with the grey-point of the ternary balance scheme. Consequently, the colors now show the direction and magnitude of the deviation from that average. Yellow, cyan, and magenta hues signify a higher-than-average share of workers in the primary, secondary, and tertiary sectors. The saturation of the colors encodes the magnitude of that deviation, with perfect grey marking a region with a workforce composition equal to the European average, i.e., the new reference point.

The centered ternary balance scheme emerges from an application of a regular ternary color-coding to transformed (centered) compositional data. In the following, I explain its construction.

3. Perturbation and centering of compositional data

Given a series of temperature readings for each day of a year, one may want to show how each reading compares to the yearly average. By subtracting the annual average from each data point, a new variable is created. It represents direction and magnitude of the temperature deviation from the mean with a reference point at zero. For the domain of compositional data, a similar transformation goes by the name perturbation, and it has been proposed as a means of centering unbalanced observations in a ternary diagram (Von Eynatten, Pawlowsky-Glahn, and Egozcue 2002; Pawlowsky-Glahn and Buccianti 2002).

When John Aitchison set out the principles of compositional data analysis (Aitchison 1982, 1986; Pawlowsky-Glahn, Egozcue, and Tolosana-Delgado 2015) he used ternary diagrams to illustrate the appropriate sample space, a positive simplex. Compositions are defined as points in the simplex or equivalently as a vector \mathbf{x} with positive elements $\langle x_1, \dots, x_D \rangle$ constrained to sum to unity. Just as the operations of linear algebra are intuitively illustrated in the two dimensions of the Cartesian plane, coordinates in the ternary diagram change in reaction to operations on the corresponding simplex, one such operation being the perturbation of a composition $\mathbf{x} = \langle x_1, \dots, x_D \rangle$ by another $\mathbf{y} = \langle y_1, \dots, y_D \rangle$ defined as $\mathbf{x} \oplus \mathbf{y} = \mathcal{C}\langle x_1 y_1, \dots, x_D y_D \rangle$, where $\mathcal{C}\langle x_1, \dots, x_D \rangle = \left\langle \frac{x_1}{\sum_i x_i}, \dots, \frac{x_D}{\sum_i x_i} \right\rangle$ denotes the closure operation that imposes the unit-sum constraint via division of each vector element by the sum of all elements. As a consequence of this definition, the perturbation of any composition \mathbf{y} by its inverse $\mathbf{y}^{-1} = \langle 1/y_1, \dots, 1/y_D \rangle$ results in the neutral element $\mathbf{y} \oplus \mathbf{y}^{-1} = \langle 1/D, \dots, 1/D \rangle$, which for a three-part composition coincides with the barycenter of a ternary diagram at $\langle 1/3, 1/3, 1/3 \rangle$. Thus, perturbation allows the transformation of coordinates in a ternary diagram such that an arbitrary composition can be relocated to the center, acting as a reference point against which all other compositions are compared.

Whenever a compositional data set is perturbed by the inverse of its compositional mean, the operation is known as centering – a key component for the construction of the centered ternary balance scheme as it allows the color-coding of the deviations from an average, thereby visualizing structure in surfaces of unbalanced data.

4. The centered ternary balance scheme


The construction of the centered ternary balance scheme is straightforward. One needs but two ingredients: 1) the ability to colorize a ternary composition with the regular ternary balance scheme, and 2) the ability to perturbate a ternary composition by the inverse of its compositional mean as proposed by Von Eynatten, Pawlowsky-Glahn, and Egozcue (2002) and described above. One first performs the centering of the compositional data set and then colorizes the centered data according to the regular ternary balance scheme. In the following, I demonstrate how these two operations resulted in the map of deviations from the average European workforce composition by region in 2016 (see Figure 2b) and propose several options for drawing an informative color key.

In 2016 the average European region had 4% of the workforce situated in the primary sector, 24% in the secondary sector, and 72% in the tertiary sector. The composition $\mathbf{y} = \langle 0.04, 0.24, 0.72 \rangle$ marks the reference against which all other compositions are to be compared.² Let $\mathbf{x}_j = \langle x_{1j}, x_{2j}, x_{3j} \rangle$ be the workforce composition of region j as shown in Figure 3a. For each region I calculate the perturbation $\mathbf{x}'_j = \mathbf{x}_j \oplus \mathbf{y}^{-1}$ shown in Figure 3b. The perturbed compositions are then color-coded according to a regular ternary balance scheme. The simplest way to achieve this is to interpret the composition as coordinates in the RGB color space. See, e.g., Wang, Robertson, and Haines (2009) for such an approach. A more flexible method employed in this paper is described in Schöley and Willekens (2017) and is implemented in the tricolore R package (Schöley and Kashnitsky 2019b). Based upon the CIE-Lch color space – a transformation of CIE-Luv described in Ware (2013), chapter 4 – the technique allows for different choices of primary colors and gives the user control over overall lightness levels and contrast while ensuring that all primary colors used in the mixing appear roughly equal in lightness and chroma. Additionally, in the CIE-Lch space, the color dimensions hue, lightness, and chroma can be manipulated truly independent of each other, a property not shared by the popular RGB or HSV color spaces (Zeileis, Hornik, and Murrell 2009).

The resulting colors show for each composition the direction and magnitude of deviation from the compositional mean. The hue of the color encodes which components of a three-part composition are greater than the average, whereas lightness and saturation indicate the distance of a composition from the average, which itself is colored grey.


² In compositional data analysis, the standard measure of centrality is the vector formed by closure of the component-wise geometric means of the observed compositions. Von Eynatten, Pawlowsky-Glahn, and Egozcue (2002) advocate for this measure to be used when centering data on the ternary diagram and it is used here as well. Note, however, that \mathbf{y} is virtually identical to the workforce distribution calculated from the numbers of workers in each sector summed across all European regions.

Figure 3: Different representations of the color key for the (centered) ternary balance scheme showing the workforce composition by region in Europe, 2016. Data by Eurostat.




A ternary diagram showing the distribution of regional European workforce compositions in 2016. The black lines mark the European average. The background is colored with the ternary balance scheme.

The data in figure a has been perturbed by the inverse of the European average. The coordinates of the resulting centered compositions are interpreted as colors in the centered ternary balance scheme.



To facilitate interpretation, the centered compositions can be labeled in terms of their original values and their percent point differences from the reference composition, here the European average, which is now situated at the barycenter of the triangle.

The centered ternary balance scheme is a skewed standard ternary balance scheme with its grey-point shifted to the location of the reference composition. Note that this color key conveys the same information as Figure c. Whereas in Figure c the data has been centered on the color scale, here the color scale has been centered on the data.



The log-ratio coordinate transformation allows us to depict a three-part composition in the unconstrained and familiar space of the real numbers. The origin of the Cartesian coordinates marks the reference composition.

There are multiple options for drawing an informative color key. Simply plotting the centered compositions in a ternary diagram with a color-coded background as in Figure 3b – while correct – is not very intuitive, as the transformed compositions cannot be easily interpreted³. A better option is to plot the centered data in a ternary diagram with scale labels indicating the original proportions (see Figure 3c). Because the centering operation skews the grid lines in a ternary diagram, centered grid lines have to be plotted (Von Eynatten, Pawlowsky-Glahn, and Egozcue 2002). This is achieved by perturbing the ternary coordinates of the grid by the inverse of the center \mathbf{y} . As the colors encode deviations from the center, the grid lines can accordingly be labeled with the percent point difference from it. A third option is to plot the uncentered data in the standard ternary diagram and perturb the background color surface instead, shifting its grey-point from the barycenter of the triangle to the location of \mathbf{y} (see Figure 3d).

A fourth legend style avoids the ternary diagram altogether and instead displays the log-ratio-transformed compositions in Cartesian coordinates (Figure 3e). This style of legend may be preferred by an audience already familiar with the methods of compositional data analysis as defined by Aitchison (1982), who introduced the so-called (additive) log-ratio transform as a means to analyze compositions in the unconstrained and familiar space of real numbers. For a three-part composition \mathbf{x} , the log-ratio transform is defined as $\text{alr}\langle x_1, x_2, x_3 \rangle = \langle \log(x_1/x_3), \log(x_2/x_3) \rangle = \langle z_1, z_2 \rangle$ with inverse $\text{alr}^{-1}\langle z_1, z_2 \rangle = \mathcal{C}\langle \exp(z_1), \exp(z_2), 1 \rangle = \langle x_1, x_2, x_3 \rangle$. The transformation maps the three positive elements of the composition to an unconstrained pair of coordinates, which may be visualized in a standard scatterplot. Notably, centering \mathbf{x} by perturbing with \mathbf{y}^{-1} is equivalent to subtracting $\text{alr}(\mathbf{y})$ from $\text{alr}(\mathbf{x})$ and applying the inverse transform to the result, illustrating the relationship between perturbation in the simplex and a simple translation of the origin in real coordinates (Aitchison and Ng 2005).

5. Discussion

With the centered ternary balance scheme – a straightforward synthesis of the three-variable balance scheme as described by Brewer (1994) and the centering operation applied in the context of compositional data analysis in the ternary diagram (Von Eynatten, Pawlowsky-Glahn, and Egozcue 2002) – I have proposed a visualization technique capable of showing the divergence of a three-part compositional surface from its average. The technique can display the internal variation of a data set, which is narrowly clustered (as demonstrated in Figure 2). More generally, the centered scheme may also be employed to show the divergence from *any* point of reference via perturbation by the inverse of the reference composition. For example, perturbing the regional distribution

³ The same situation arises when a logarithmic scale is labeled with the logged values as opposed to the values on the original scale.

of educational attainment in Figure 1b by the average composition in the U.S. would yield a map of compositional differences from the U.S. average. Further, compositional change over time can be visualized by perturbing each data point at t_2 by the inverse of the corresponding composition at t_1 (Aitchison and Ng 2005) and colorizing the resulting perturbation using the standard ternary balance scheme.

Neither the ternary balance scheme nor its centered extension has been empirically tested regarding the effectiveness of visualizing compositional data on a surface. However, visualization theory gives some insight into potential strengths and weaknesses of the technique. The ternary balance technique uses the visual attribute “color” as a multidimensional encoding for the three parts that make up a ternary composition, mapping each part of the composition to a separate primary color channel. Is it possible for a reader of the visualization to separate the ternary colors into their three primaries, thereby perceiving the relative magnitude of each compositional part separately? The answer is no. Color primaries are known to be integral: they are perceived jointly and are hard to separate (Ware 2013). Therefore, the ternary balance technique should not be used in situations where it is essential to precisely identify the relative shares of three components. However, the scheme can also be interpreted as a hue-(lightness/chroma) encoding. While hue signifies a qualitative attribute – the dominant part(s) of the composition – lightness and chroma redundantly encode the distance of a compositional observation from a reference composition – a quantitative attribute. Hue and lightness can be separated to some degree, as illustrated by color names such as light blue, dark green, and so forth. Interpreting the hues and lightness components of a map colored with the ternary balance scheme allows numerous relevant tasks to be performed:

1. **Identification of regions close to the reference composition.** Locate dark and grey regions.
2. **Identification of regions deviating from the reference composition.** Locate bright and colorful regions.
3. **Classification of regions deviating from the reference composition.** Identify the hue of the region and associate the hue with its corresponding part(s) of the composition.
4. **Identification of compositional spatial gradients.** Locate color gradients.
5. **Classification of compositional spatial gradients.** Identify the starting hue and ending hue of the gradient. Associate the starting and ending hues with their corresponding part of the composition.
6. **Identification of compositional spatial discontinuities.** Locate sudden shifts in hue and lightness.

- 7. Classification of compositional spatial discontinuities.** Identify the hues at both sides of the discontinuity and associate them with the corresponding part of the composition.

None of the above tasks requires the reader to perform the impossible feat of decomposing a color into its primary constituents. The identification tasks 1 and 2 can be performed solely by comparing lightness levels. Identification tasks 4 and 6 require judging the dissimilarity of neighboring colors. Classification tasks 3, 5, and 7 require the rough matching of two hues. All of these perception and cognitive tasks can also be found in established visualization techniques. Therefore I hypothesize that they can also be performed on the ternary balance scheme. Empirical evidence has to be collected to support that claim.


Whenever hue is used as a visual encoding, consideration has to be given to people with impaired color vision. Because the ternary balance scheme relies on the mixture of three distinct primaries, the resulting space of possible colors will necessarily contain colors that are non-discriminable by people with impaired color vision. The most common form of color-blindness reduces the sensitivity to green light, making it hard to distinguish colors along the red-green spectrum (Birch 2012). If the compositional data set does not cover the whole surface of the ternary legend, then it may be possible to choose the primary colors such that the data fall outside of the red-green spectrum.

The technique described in this paper is implemented in the `tricolore` R package (Schöley and Kashnitsky 2019b),⁴ which builds upon the facilities of the `ggtern` package (Hamilton and Ferry 2018) to draw color-coded ternary diagrams. Given a three-column matrix of three-part compositions, `tricolore` returns a vector of colors and a suitable color key. The user may choose between discrete and continuous color scales, center the color scale around either the geometric mean of the provided data or an arbitrary reference composition, and experiment with different values for hue, lightness, chroma, and contrast. An online tutorial (Schöley 2019) explains how to use the software to create color-coded maps like those shown in this paper, and a user interface (see Figure 4) helps with picking suitable parameters for the color scale.

⁴ Development takes place in a public repository at <https://github.com/jschoeley/tricolore> and the authors welcome suggestions and bug reports.

Figure 4: The tricolore package for the statistical programming language R implements the centered ternary balance color scheme and provides a user interface for quickly testing different parametrizations.

Tricolore: A flexible color scale for ternary compositions



Thus far the tricolore software has been used in publications to map regional deviations from the average European age composition (Kashnitsky and Schöley 2018; Schöley and Kashnitsky 2019a), disparities in regional education attainment in India and Nigeria (Graetz et al. 2019), the cause of death distribution by period, age, and region in Mexico (Kashnitsky and Aburto 2019), and Vienna's district population mix by region of origin (Stadt Wien 2019). I hope that tricolore continues to encourage people to experiment with this novel visualization technique – three-part compositions are plenty, and surfaces, whether defined by longitude and latitude or by period and age, provide ample room to find exciting variation in the data.

6. Acknowledgments

My thanks go to the three reviewers who generously contributed their time and expertise. I would also like to thank Marie-Pier Bergeron Boucher and Jim Oeppen for their input regarding the analysis of compositional data and Ilya Kashnitsky, who inspired me to write this paper.

References

- Aitchison, J. and Ng, K.W. (2005). The role of perturbation in compositional data analysis. *Statistical Modelling* 5(2): 173–185. doi:10.1191/1471082X05st091oa.
- Aitchison, J. (1982). The statistical analysis of compositional data. *Journal of the Royal Statistical Society. Series B* 44(2): 139–177. doi:10.2307/2347798.
- Aitchison, J. (1986). *The statistical analysis of compositional data*. Monographs on statistics and applied probability. New York: Chapman and Hall.
- Bancroft, W.D. (1897). A triangular diagram. *Journal of Physical Chemistry* 1(7): 403–410. doi:10.1021/j150589a002.
- Birch, J. (2012). Worldwide prevalence of red-green color deficiency. *Journal of the Optical Society of America A* 29(3): 313–320. doi:10.1364/JOSAA.29.000313.
- Brewer, C.A. (1994). Color use guidelines for mapping and visualization. In: MacEachren, A.M. and Taylor, D.R.F. (eds.). *Visualization in Modern Cartography*. Oxford, UK: Pergamon: chap. 7: 123–147, Modern Cartography. doi:10.1016/b978-0-08-042415-6.50014-4.
- Davis, R.O.E. and Hammond, B.H. (1927). *Grouping of soils on the basis of mechanical analysis*. Department Circular 419, Washington D.C. <https://archive.org/details/groupingofsoilso419davi>.
- De Finetti, B. (1926). Considerazioni matematiche sull'ereditarietà mendeliana. *METRON* VI(1): 3–411.
- Denil, M. (2015). Trivariate sea ice presence map. In: *6th Symposium on the Impacts of an Ice-Diminishing Arctic on Naval and Maritime Operations*. National Ice Center. https://www.star.nesdis.noaa.gov/star/documents/meetings/Ice2015/posters/Denil_M_poster.png.
- Dorling, D. (2012). *The visualization of spatial social structure*. Wiley Series in Computational and Quantitative Social Science. Chichester, UK: Wiley. doi:10.1002/9781118353929.
- Graetz, N., Woyczynski, L., Wilson, K.F., Hall, J.B., Abate, K.H., Abd-Allah, F., Adebayo, O.M., Adekanmbi, V., Afshari, M., Ajumobi, O. et al. (2019). Mapping disparities in education across low-and middle-income countries. *Nature* 577: 235–238. doi:10.1038/s41586-019-1872-1.
- Hamilton, N.E. and Ferry, M. (2018). ggtern: Ternary diagrams using ggplot2. *Journal of Statistical Software* 87(1): 1–17. doi:10.18637/jss.v087.c03.

- Howarth, R.J. (1996). Sources for a history of the ternary diagram. *The British Journal for the History of Science* 29(3): 337–356. doi:10.1017/S000708740003449X.
- Kashnitsky, I. and Aburto, J.M. (2019). Geofaceting: Aligning small-multiples for regions in a spatially meaningful way. *Demographic Research* 41(17): 477–490. doi:10.4054/DemRes.2019.41.17.
- Kashnitsky, I. and Schöley, J. (2018). Regional population structures at a glance. *The Lancet* 392(10143): 209–210. doi:10.1016/s0140-6736(18)31194-2.
- Kvitkin, O.A. (1932). *On types of urban settlements [o tipakh gorodskikh poseleniy]*. USSR census 1926.
- Metternicht, G. and Stott, J. (2003). Trivariate spectral encoding: a prototype system for automated selection of colours for soil maps based on soil textural composition. In: *Proceedings of the 21st International Cartographic Conference (ICC) "Cartographic Renaissance"*. 2341–2353.
- Pawlowsky-Glahn, V. and Buccianti, A. (2002). Visualization and modeling of sub-populations of compositional data: statistical methods illustrated by means of geochemical data from fumarolic fluids. *International Journal of Earth Sciences* 91(2): 357–368. doi:10.1007/s005310100222.
- Pawlowsky-Glahn, V., Egozcue, J.J., and Tolosana-Delgado, R. (2015). *Modeling and analysis of compositional data*. Statistics in practice. United Kingdom: John Wiley & Sons. doi:10.1002/9781119003144.
- Pirzamanbein, B., Poska, A., and Lindström, J. (2020). Bayesian reconstruction of past land cover from pollen data: Model robustness and sensitivity to auxiliary variables. *Earth and Space Science* 7: 1–13. doi:doi.org/10.1029/2018EA000547.
- Schöley, J. (2019). Choropleth maps with tricolore. online. https://cran.r-project.org/web/packages/tricolore/vignettes/choropleth_maps_with_tricolore.html.
- Schöley, J. and Kashnitsky, I. (2019a). But Why? Design choices made while creating "Regional population structures at a glance". In: Fischer, J., Mazouch, P., Tesárková, K.H., and Kurtinová, O. (eds.). *New Generations in Demography*. Oeconomica Publishing House. doi:10.18267/pu.2019.fis.2302.6.
- Schöley, J. and Kashnitsky, I. (2019b). tricolore: A flexible color scale for ternary compositions. CRAN. <https://cran.r-project.org/package=tricolore>. V1.2.1.
- Schöley, J. and Willekens, F. (2017). Visualizing compositional data on the Lexis surface. *Demographic Research* 36(21): 627–658. doi:10.4054/DemRes.2017.36.21.
- Stadt Wien (2019). Wien in Zahlen 1919–1934. <https://www.wien.gv.at/statistik/pdf/rotes-wien-bildband.pdf>.

- Steidinger, B.S., Crowther, T.W., Liang, J., Van Nuland, M.E., Werner, G.D.A., Reich, P.B., Nabuurs, G.J., de Miguel, S., Zhou, M., Picard, N. et al. (2019). Climatic controls of decomposition drive the global biogeography of forest-tree symbioses. *Nature* 569(7756): 404–408. doi:doi.org/10.1038/s41586-019-1128-0.
- Von Eynatten, H., Pawlowsky-Glahn, V., and Egozcue, J.J. (2002). Understanding perturbation on the simplex: A simple method to better visualize and interpret compositional data in ternary diagrams. *Mathematical Geology* 34(3): 249–257. doi:[10.1023/A:1014826205533](https://doi.org/10.1023/A:1014826205533).
- Wang, Q.J., Robertson, D.E., and Haines, C.L. (2009). A bayesian network approach to knowledge integration and representation of farm irrigation. *Water Resources Research* 45(2). doi:[10.1029/2006WR005421](https://doi.org/10.1029/2006WR005421).
- Ware, C. (2013). *Information visualization. Perception for design*. Waltham, MA: Elsevier, 3 ed. doi:[10.1016/C2009-0-62432-6](https://doi.org/10.1016/C2009-0-62432-6).
- Zabetakis, M.G. (1965). *Flammability characteristics of combustible gases and vapors*. Tech. Rep. 627, Washington D. C. <https://apps.dtic.mil/dtic/tr/fulltext/u2/701576.pdf>.
- Zeileis, A., Hornik, K., and Murrell, P. (2009). Escaping RGBland: Selecting colors for statistical graphics. *Computational Statistics & Data Analysis* 53(9): 3259–3270. doi:[10.1016/j.csda.2008.11.033](https://doi.org/10.1016/j.csda.2008.11.033).

